

学術情報リポジトリ

Ketoconazole-induced estrogen deficiency causes transient decrease in placental blood flow associated with hypoxia and later placental weight gain in rats

メタデータ	言語: eng
	出版者:
	公開日: 2017-08-31
	キーワード (Ja):
	キーワード (En):
	作成者: Ichikawa, Aoi, Tamada, Hiromichi
	メールアドレス:
	所属:
URL	http://hdl.handle.net/10466/15560

		Treatment period of KTZ (day of pregnancy)		
	Control	9-11	12-14	15-17
No. of dams	6	5	7	7
No. of live fetuses per dam	$11.5\pm2.5$	$13.8\pm0.8$	$13.0\pm1.7$	$14.3\pm1.8$
Fetal weight (g)				
Male	$3.57\pm0.34$	$3.49\pm0.17$	$3.60\pm0.33$	$3.63\pm0.24$
Female	$3.36\pm0.22$	$3.35\pm0.21$	$3.42\pm0.26$	$3.48\pm0.25$
Placental weight (mg)				
Male	$492\pm59^{a}$	$458\pm18^{a}$	$626\pm88^{b}$	$516 \pm 33^{a}$
Female	$476\pm54^a$	$433\pm36^a$	$614\pm73^b$	$492 \pm 49^{a}$

## Table 1Effects of KTZ treatment during various periods of pregnancy on the weight of fetuses and placentas.

Daily oral administration of KTZ (25 mg/kg) was given on Days 9 to 11, 12 to 14 or 15 to 17 of pregnancy and the fetuses and placentas were weighed on Day 20. Values are means  $\pm$  SEM. <sup>ab</sup>Values of placental weights with different letters differ significantly at *p*<0.05. There is no difference in the placental weights between male and female fetuses, while the values in the rats treated with KTZ on Days 12 to 14 are greater than the other groups regardless of fetal gender.

## Table 2

Effects of KTZ and  $E_2$  on percentage of sinusoid area per unit area in the labyrinth zone of placentas on Day 20 of pregnancy.

	Control	KTZ+0E2	KTZ+0.1E2	KTZ+1E <sub>2</sub>
Sinusoid area (%)	$19.15 \pm 1.63^{a}$	$46.16 \pm 2.32^{\circ}$	$29.23\pm2.93^{b}$	$27.62\pm2.92^{ab}$

KTZ (25 mg/kg/day) and E<sub>2</sub> (0, 0.1 or 1  $\mu$ g/rat/day) were treated on Days 12 to 14 of pregnancy. Values are mean  $\pm$  SEM (n=6). <sup>abc</sup>Values with different letters differ significantly at *p*<0.05. The value in the KTZ+0E<sub>2</sub> group is greater than those in the other groups, and the value in the KTZ+0.1E<sub>2</sub> group is larger than that in controls.

Table 3

Effects of KTZ and  $E_2$  on the number of erythrocytes and sinusoid area per unit area in the labyrinth zone of placentas on Day 14 of pregnancy.

	Control	KTZ+0E <sub>2</sub>	KTZ+1E <sub>2</sub>
No. of all erythrocytes	831.5 ± 75.5	$838.0 \pm 119.5$	$1100.0 \pm 102.5$
Ratio of fetal erythrocyte (%)	$17.7 \pm 4.0^{a}$	$45.2\pm3.8^{\rm b}$	$14.8\pm2.8^{\rm a}$
Maternal sinusoid per unit area (%)	$20.2\pm3.2^{\rm a}$	$11.9\pm1.8^{\rm b}$	$31.6 \pm 1.4^{c}$
Fetal sinusoid per unit area (%)	$4.6\pm0.6^{\rm a}$	$10.8\pm0.6^{\rm b}$	$5.8\pm0.9^{\mathrm{a}}$

KTZ (25 mg/kg/day) and E<sub>2</sub> (0 or 1  $\mu$ g/rat/day) were administered on Days 12 to 14 of pregnancy. Values are mean  $\pm$  SEM (n=6). <sup>abc</sup>Values with different letters within each row differ significantly at p<0.05.