



Examples of Algebraic Varieties with Kobayashi Hyperbolicity

メタデータ	言語: eng 出版者: 公開日: 2013-12-19 キーワード (Ja): キーワード (En): 作成者: Matsuno, Takanori メールアドレス: 所属:
URL	https://doi.org/10.24729/00007623

EXAMPLES OF ALGEBRAIC VARIETIES WITH KOBAYASHI HYPERBOLICITY

Takanori MATSUNO*

Abstract

Many interesting examples of hyperbolic hypersurfaces in the complex projective space $\mathbf{P}^3(\mathbf{C})$ have been known. In this paper, we give some examples of 2-dimensional hyperbolic algebraic varieties which are defined as intersections of Fermat varieties in $\mathbf{P}^4(\mathbf{C})$.

Key Words : Kobayashi hyperbolicity, Nevanlinna theory

1 Introduction

In [K], Kobayashi asked whether a generic hypersurface in the complex projective space $\mathbf{P}^n(\mathbf{C})$ of degree enough large with respect to n is hyperbolic or not. This conjecture is true for $n = 2$. In fact, a plane curve with genus greater than or equal to 2, does not admit any non-constant holomorphic mapping from \mathbf{C} , because its universal covering space is a ball. For $n \geq 3$ this problem is still open. But there have been known many examples of hyperbolic hypersurfaces in $\mathbf{P}^3(\mathbf{C})$ ([Br-Gr][D][F2][Go][N][S1][S2]). In this paper, we give some examples of 2-dimensional hyperbolic algebraic varieties which are defined as intersections of Fermat varieties in $\mathbf{P}^4(\mathbf{C})$.

2 Preliminaries

We recall some definitions and a result.

Definition 2.1. For two entire functions f and g which are not identically zero, we say they are equivalent if there exists a constant c ($c \neq 0$) such as $f = cg$ holds. This introduces an equivalence relation in each set of entire functions which are not identically zero. We mean by the notation $f \sim g$ that f and g are equivalent.

Definition 2.2. Let f be a holomorphic mapping of \mathbf{C} into $\mathbf{P}^n(\mathbf{C})$. A representation $\tilde{f} = (f_0, \dots, f_n)$ of f is a holomorphic mapping of \mathbf{C} into \mathbf{C}^{n+1} such that $\tilde{f}^{-1}(\mathbf{0}) \neq \mathbf{C}$ and $f(z) = (f_0(z) : \dots : f_n(z))$ for each $z \in \mathbf{C} \setminus \tilde{f}^{-1}(\mathbf{0})$, where $(X_0 : \dots : X_n)$ is a homogeneous coordinate system. A representation \tilde{f} is called to be reduced if $\tilde{f}^{-1}(\mathbf{0}) = \emptyset$.

The following theorem was given by Green [Gr] and Fujimoto [F1]:

Theorem 2.1. Let f_0, \dots, f_n be entire functions which are not identically zero such that $f_0^d + \dots + f_n^d \equiv 0$, where d is a positive integer. If $d \geq n^2$, then

$$\sum_{f_j \in I} f_j^d \equiv 0$$

for each equivalence class I of $\{f_0, \dots, f_n\}$. Especially each class has at least two elements.

3 Main theorem

Let d be a positive integer. Put $M_d := \{X_0^d + \dots + X_4^d = 0\}$, where X_0, \dots, X_4 are homogeneous coordinates, which is a Fermat variety of degree d in $\mathbf{P}^4(\mathbf{C})$. First take d as d is greater than or equal to 16. And then take d' such as the set of d -th roots of -1 and that of the d' -th roots of -1 do not share any element. We define a complex surface S in $\mathbf{P}^4(\mathbf{C})$ as $S := M_d \cap M_{d'}$. Then we have the following theorem:

Theorem 3.1. S is Kobayashi hyperbolic.

Proof. Assume that there exists a holomorphic mapping f of \mathbf{C} into $\mathbf{P}^4(\mathbf{C})$ with reduced representation $\tilde{f} = (f_0, f_1, f_2, f_3, f_4)$ such that $f(\mathbf{C}) \subset S$. Since $f(\mathbf{C}) \subset M_d$, $f_0^d + \dots + f_4^d \equiv 0$.

First we assume that each f_j is not identically zero. By **Theorem 2.1**, the set $\{f_0, \dots, f_4\}$ of entire functions can be divided into each equivalence classes. Let N be the number of elements of an equivalence class of f_0 .

(Received April 11, 2007)

* Dept. of Industrial Systems Engineering : Natural Science

(I) The case that $N = 5$. Clearly, f is constant in this case.

(II) The case that $N = 4$. This case cannot occur. Because each equivalent class has at least two elements.

(III) The case that $N = 3$. In this case, say, $f_1 = c_1 f_0$, $f_2 = c_2 f_0$, $f_3 = c_3 f_4$. By Theorem 2.1, we get $1 + c_1^d + c_2^d = 0$ and $c_3^d + 1 = 0$. But by the assumption for d' , this does not satisfy the condition that $f(\mathbf{C}) \subset M_{d'}$. Then this case cannot occur.

(IV) The case that $N = 2$. This case is as same as the above. Then this case cannot occur.

(V) The case that $N = 1$. This case cannot occur. Because each equivalent class has at least two elements.

Next we consider the case that $f_j \equiv 0$ for some j . In the case $f_j \equiv 0$ for only one j , the each equivalence class has 4 or 2 elements. If an equivalence class has 4 elements, it is clear that f is constant. If each equivalence class has 2 elements, the condition that $f(\mathbf{C}) \subset M_{d'}$ is not satisfied, same as above (III). Then this case cannot occur. In the case that $f_j \equiv 0$ for more than 2 j 's, it is clear that f is constant since the image $f(\mathbf{C})$ is included in a hyperbolic Riemann surface.

We have shown that every holomorphic mapping f of \mathbf{C} into S is constant. So S is hyperbolic. The proof is completed.

For example, if we take d such as d is even and is greater than or equal to 16 and put $d' = 1$, then S is biholomorphic to the hypersurface $S_d := \{X_0^d + X_1^d + X_2^d + X_3^d + (-X_0 - X_1 - X_2 - X_3)^d = 0\}$ in $\mathbf{P}^3(\mathbf{C})$. Then we have:

Theorem 3.2. S_d is a hyperbolic hypersurface in $\mathbf{P}^3(\mathbf{C})$.

Remark. This example of a hyperbolic hypersurface is not a new one. This is given in a rather complicated situation in [S1].

References

- [Br-Gr] R. Brody and M. Green : *A family of smooth hyperbolic hypersurfaces in \mathbf{P}^3* , Duke Math. J., 44(1977), 873-874
- [D] J. P. Demailly : *Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials*, Proc. Sympos. Pure Math., Vol. 62, Part2, Amer. Math. Soc., Providence, RI, 1997, 285-360.
- [F1] H. Fujimoto : *On meromorphic maps into the complex projective space*, J. Math. Soc. Japan, 26(1974), 272-288.
- [F2] H. Fujimoto : *A family of hyperbolic hypersurfaces in the complex projective space*, The Chuang special issue. Complex Variables Theory Appl., 43(2001), no. 3-4, 273-283.
- [Go] J. Goul : *Algebraic families of smooth hyperbolic hypersurfaces of low degree in $\mathbf{P}_{\mathbf{C}}^3$* , manuscripta math., 90(1996), 521-532.
- [Gr] M. L. Green : *Some Picard theorems for holomorphic maps to algebraic varieties*, Amer. J. Math., 97(1975), 43-75.
- [K] S. Kobayashi : *Hyperbolic Manifolds and Holomorphic Mappings*, Dekker, 1987.
- [N] A. Nadel : *Hyperbolic surfaces in \mathbf{P}^3* , Duke Math. J., 58(1989), 749-771
- [S1] M. Shirotsaki : *Hyperbolic hypersurfaces in the complex projective spaces of low dimensions*, Kodai Math. J., 23(2000), 224-233.
- [S2] M. Shirotsaki : *A hyperbolic hypersurface of degree 10*, Kodai Math. J., 23(2000), 376-379.